

A category-theoretic reformulation of Shelah's dividing lines in model theory.

P&S'60

During poster room hours at P&S'60:
chat: t.me/McVlr
voice: mishap.sdf.org/gathertown

Misha Gavrilovich (IPRERAS), Konstantin Pimenov (St.Petersburg State University).

Thanks due to M.Bays and A.Hasson.

1. Summary

We interpret category-theoretically several of the Shelah dividing lines in model theory, using a simplicial category of generalised topological spaces. This makes formal the intuition that these properties of models and formulas are defined in terms of avoiding certain “bad” infinite combinatorial structures: the same diagram chasing “trick”, the lifting property, applied to (a morphism associated with) a combinatorial structure defines the associated no-tree- or no-order- property of (objects associated with) models. The list of properties includes NOP, NTP, NATP, NTP_i , $NSOP_i$ ($i \geq 1$) and NIP.

A logician should find amusing a metamathematical aspect—**how trivial is all we do here: take the text of the usual definition of NTP in Tent-Ziegler [5] and “transcribe” it line by line into the simplicial language** in an *oversimplified* manner. [9] does the same with a few other textbook definitions including that of a topological space and a limit of a filter in (Bourbaki, General Topology).

2. A category of generalised topological spaces

We define our generalised topological spaces to be simplicial objects of the category of filters on sets, or, equivalently, the category of finitely additive measures taking values 0 and 1 only. Thus a **generalised topological space is a simplicial set equipped, for each $n \geq 0$, with a filter on the set of n -simplices such that under any face or degeneration map the preimage of a large set is large.**

These spaces generalise uniform and topological spaces, filters, and simplicial sets, and the concept is designed to be flexible enough to formulate categorically a number of standard basic elementary definitions in various fields, e.g. in analysis, limit, (uniform) continuity and convergence, equicontinuity of sequences of functions; in algebraic topology, being locally trivial and geometric realisation; in geometry, quasi-isomorphism; in model theory, stability, simplicity and several Shelah's dividing lines [3,9].

3. The definition of generalised topological spaces

Definition (Continuous maps of filters). *Let X and Y be sets equipped with filters (resp. measures). Call a map $f : X \rightarrow Y$ continuous iff the preimage of a big (resp. full measure) set is necessary big (resp. has full measure).*

Definition (Generalised topological spaces). *Let \mathcal{F} denote the category formed by sets equipped with filters, and their continuous maps. Its category of simplicial objects*

$$s\mathcal{F} := \text{Functors}(\text{Non-emptyFiniteLinearOrders}^{\text{op}}, \mathcal{F})$$

is our category of generalised topological spaces.

In $s\mathcal{F}$ a topological, resp. uniform, space X is the simplicial set represented by the set of points of X , where $X \times X$ is equipped with the filter of non-uniform neighbourhoods of the diagonal of form $\bigsqcup_{U_x} \text{a neighbourhood of } x \in X \{x\} \times U_x$, resp. the uniformity filter, and each X^n is equipped with the coarsest filter such that all the simplicial maps $X^n \rightarrow X \times X$ are continuous. Geometric realisation is a space of maps in $s\mathcal{F}$ from $[0, 1]^{\mathbb{N}}$ to a simplicial set [3,8,9].

4. A precise meaning for “ n -tuple being sufficiently small” for $n > 2$

A topological structure on a set enables one to give an exact meaning to the phrase “whenever x is sufficiently near a , x has the property $P(x)$ ”, whereas a **generalised topological space enables one to give an exact meaning to the phrase “every n -tuple of sufficiently similar points x_1, x_2, \dots, x_n has property $P(x_1, \dots, x_n)$ ” for $n > 1$.** (Uniform spaces were introduced to do this for $n = 2$ and “similar” meaning “at small distance”, as explained in (Bourbaki, General Topology; Introduction)). In a topological space, this exact meaning is that the set $\{x \mid P(x)\}$ belongs to the neighbourhood filter of a point a . Similarly, in a generalised topological space, it is that the set $\{(x_1, \dots, x_n) \mid P(x_1, \dots, x_n)\}$ belongs to the “neighbourhood” filter defined on n -simplices.

In model theoretic examples, similarity may mean either indiscernability or realising sufficiently many instances of a formula: n -simplices are tuples of elements of a model, and the “neighbourhood filter” on n -tuples consists of all subsets containing all “sufficiently” indiscernible tuples or realising “sufficiently many” instances of a formula. **This exact meaning enables us to bring the standard intuition of topology to model theory.**

5. The definition of NTP (no tree property)

For a binary formula $\varphi(x, y)$, call a tuple $(a_i : 0 < i < n)$ φ -consistent iff the set $(\varphi(x, a_i) \mid 0 < i < n)$ is consistent. Below ${}^{<\omega}\omega$, resp. ${}^{\omega}\omega$, denotes the tree formed by finite, resp. countably infinite, sequences of natural numbers.

Definition (Tent-Ziegler, 7.2.1). *A formula $\varphi(x, y)$ has the tree property (with respect to k) if there is a tree of parameters $(a_s \mid \emptyset \neq s \in {}^{<\omega}\omega)$ such that:*

- For all $s \in {}^{<\omega}\omega$, each k -tuple of distinct siblings $(a_{s_i} \mid i < k)$ is φ -inconsistent.
- For all branches $\sigma \in {}^{\omega}\omega$ the tuple $(a_s \mid \emptyset \neq s \subseteq \sigma)$ is φ -consistent.

What do you gain by pretending so ?

[Read the definition of NTP line by line and rephrase it in terms of generalised topological spaces](#)

6. The space associated with a formula

The definition cares about φ -consistency of tuples in a model M . Therefore we consider the simplicial set $M_{\bullet} : \text{Non-emptyFiniteLinearOrders}^{\text{op}} \rightarrow \text{Sets}$

$$M_{\bullet}(n^{\leq}) := \text{Hom}_{\text{sets}}(n^{\leq}, |M|) = |M|^n, n > 0$$

and equip each $|M|^n$ with the filter generated by a single subset:

the subset of φ -consistent n -tuples.

7. Rewriting the positive requirement of item b)

View a tree as a partial order under prefix relation.

The definition also cares about branches of the tree $T := {}^{<\omega}\omega$, i.e. sets of increasing tuples

$$T_{\bullet}^{\leq}(n^{\leq}) := \text{Hom}_{\text{preorders}}(n^{\leq}, T^{\leq}), n > 0.$$

An easy simplicial argument (as both are representable) shows that

a tree of parameters $(a_s \mid \emptyset \neq s \in T)$ is the same as a map $T_{\bullet}^{\leq} \rightarrow M_{\bullet}$.

We want **this map to be continuous iff the tree of parameters satisfies the positive requirement on a witness of the tree property**, i.e. item b). By definition, this means that the subset of φ -consistent tuples in each $|M|^n$ contains all the linearly ordered (in the tree) tuples of parameters. This becomes precisely what is required by the definition of continuity if we

equip each $T_{\bullet}^{\leq}(n^{\leq})$ with the indiscrete filter.

8. Rewriting the negative requirement of item a)

The negative requirement of the definition (i.e. item a)) cares about unordered tuples of parameters. To capture arbitrary tuples, consider the simplicial set

$$|T|_{\bullet}(n^{\leq}) := \text{Hom}_{\text{sets}}(n^{\leq}, |T|) = |T|^n, n > 0$$

We want **the map $|T|_{\bullet} \rightarrow M_{\bullet}$ to be continuous iff there is a witness of failure of the negative requirement (i.e. a φ -consistent k -tuple of distinct siblings) in each subtree of shape required by the definition (i.e. a copy of ${}^{<\omega}\omega$).** We turn this requirement into a definition of filters on $|T|_{\bullet}$: a subset of $|T|_{\bullet}(n^{\leq})$ is **big** iff **it contains a tuple with k distinct siblings (in the subtree) from each subtree of T isomorphic to ${}^{<\omega}\omega$, and is closed under permutations.**

A Ramsey theory argument verifies this indeed defines a filter. (A technicality: we need to slightly modify these filters to ensure $T_{\bullet}^{\leq} \rightarrow |T|_{\bullet}$ is continuous, and insist that big subsets also contain all the linearly ordered tuples of vertices.)

9. Finally, a reformulation via the lifting property

Finally, we see that

a continuous map $T_{\bullet}^{\leq} \rightarrow M_{\bullet}$ extends continuously to $|T|_{\bullet} \rightarrow T_{\bullet}^{\leq}$

iff

the tree of parameters $(a_s \mid \emptyset \neq s \in T)$ satisfying item b), has a subtree of shape ${}^{<\omega}\omega$ failing item a).

The latter is equivalent to failure of the tree property when quantified over a_i 's; the former is represented by a commutative diagram known as *lifting property*:

$$\begin{array}{ccc} T_{\bullet}^{\leq} & \xrightarrow{\quad \vee \quad} & M_{\bullet} \\ \downarrow & \nearrow & \downarrow \\ |T|_{\bullet} & \xrightarrow{\quad \vee \quad} & T \end{array}$$

This is our category-theoretic interpretation of NTP (no tree property). The same method leads to similar diagrams defining NTP_i , NATP, $NSOP_1$ (new item a) leads to new filters), and NOP, $NSOP_i$, $i \geq 3$ (also need to modify M_{\bullet}).

10. Research directions

- The lifting property is used to axiomatise homotopy theory as the notion of a Quillen model category. **Is there a homotopy theoretic interpretation of the Shelah dividing lines in model theory ?** Say, what is the model theoretic meaning of the number of connected components of the space of maps from T_{\bullet}^{\leq} to M_{\bullet} where, perhaps, the space of maps is as defined in the $s\mathcal{F}$ reformulation of geometric realisation ?
- Pillay-Starchenko [1, Cor.1.2] uses φ -consistent tuples (take $\varphi := E$ there), i.e. data captured by M_{\bullet} , and so does Simon [2, Def.1.1] to define generically stable measures. Malliaris [6] studies Szemerédi regularity of M_{\bullet} viewed as a multigraph. Reformulate this in terms of $s\mathcal{F}$. **Can definable and Szemerédi regularity be reformulated in terms of our generalised spaces ?**
- Reformulating in $s\mathcal{F}$ the Bourbaki definition of a limit of a filter on a topological space defines a notion of **limit of an arbitrary morphism in $s\mathcal{F}$ reminiscent of homotopy**, see §23. A combinatorial structure on a model, say an NTP tree, is a morphism in $s\mathcal{F}$, hence there is a notion of **a limit of a combinatorial structure on a model**. Is it useful? Can one **define homotopy of models**?

11. Appendix: The lifting property

Definition. A morphism $A \xrightarrow{f} B$ in a category has the *left lifting property with respect to a morphism* $C \xrightarrow{g} D$ and g also has the *right lifting property with respect to* f , denoted $f \perp g$, iff for each map $A \xrightarrow{t} C$ and $B \xrightarrow{b} D$ such that $f \circ b = t \circ g$, there exists $B \xrightarrow{d} C$ such that $t = f \circ d$ and $b = d \circ g$.

For a class C of morphisms in a category, its left orthogonal $C^{\perp l}$ with respect to the lifting property, respectively its right orthogonal $C^{\perp r}$, is the class of all morphisms which have the left, respectively right, lifting property with respect to each morphism in the class C .

It is clear that $C^{\perp l r} \supset C$, $C^{\perp r l} \supset C$, $C^{\perp l} = C^{\perp l r l}$, and $C^{\perp r} = C^{\perp r l r}$. The class $C^{\perp r}$ is always closed under retracts, pullbacks, (small) products (whenever they exist in the category) and composition of morphisms, and contains all isomorphisms. Meanwhile, $C^{\perp l}$ is closed under retracts, pushouts, (small) coproducts and transfinite composition (filtered colimits) of morphisms (whenever they exist in the category), and also contains all isomorphisms. A desirable property, sometimes proved by a Quillen small object argument, is that each morphism f decomposes as $f = f_l \circ f_r$ and $f = f_{rl} \circ f_r$ where $f_l \in C^{\perp l}$, $f_r \in C^{\perp l r}$, $f_{rl} \in C^{\perp r}$, and $f_{rl} \in C^{\perp r l}$.

Taking the orthogonal ("negation") of a class C is a simple way to define a class of morphisms excluding non-isomorphisms from C , in a way which is useful in a diagram chasing computation. A useful intuition is to think that the property of left-lifting against a class C is a kind of negation of the property of being in C , and that right-lifting is also a kind of negation, and for this reason it is convenient to say that

property $C^{\perp l}$, resp. $C^{\perp r}$, is left, resp. right, Quillen negation of property C .

12. Appendix: Examples of the lifting property

Iterated lifting property (=iterated Quillen negation) is used to define a number of textbook notions starting from an explicitly given list of (counter)examples or a simple property, see [4] for a list. We sketch the simplest examples.

Example. It is easy to verify that

- A map is surjective iff it has the right lifting property with respect to the map from the emptyset to a singleton (the archetypal example of non-surjection $\emptyset \rightarrow \{\bullet\}$), and it is injective iff it has the right lifting property with respect to the map gluing together two points (the archetypal non-injection example of $\{\bullet, \bullet\} \rightarrow \{\bullet\}$)

These statements are represented by the following commutative diagrams:

$$\begin{array}{ccc} \emptyset & \longrightarrow & X \\ \downarrow & \dashrightarrow & \downarrow \\ \{\bullet\} & \longrightarrow & Y \end{array} \quad \begin{array}{ccc} \{\bullet, \bullet\} & \longrightarrow & X \\ \downarrow & \dashrightarrow & \downarrow \\ \{\bullet\} & \longrightarrow & Y \end{array}$$

(surj) (inj)

where \dashrightarrow marks the property being defined.

13. Rewriting combinatorial characterisations of other dividing lines

Let us now clarify what we said earlier: The same method leads to the same diagram defining NTP_i, NATP, NSOP₁ (new item a) leads to new filters), and NOP, NSOP_i, $i \geq 3$ (also need to modify M_\bullet). We also observe that NFCP (no finite cover property) means "of infinite simplicial dimension".

14. Rewriting NTP₁ (no tree property)

Let us show the same method leads us to rewrite NTP₁ in $s\mathcal{P}$. The cases of NATP and NSOP₁ are similar.

Definition. A formula $\varphi(x, y)$ has TP₁ (tree property 1) iff there is a tree of parameters $(a_s \mid \emptyset \neq s \in {}^{<\omega}\omega)$ such that:

- for each pair $\eta, \mu \in {}^{<\omega}\omega$ of incomparable elements, the pair $\{a_\mu, a_\eta\}$ is φ -inconsistent.
- For all branches $\sigma \in {}^{<\omega}\omega$ the tuple $(a_s \mid \emptyset \neq s \subseteq \sigma)$ is φ -consistent.

Note that the only difference from the definition of NTP is in item a). Therefore, the only the only difference from the reformulation of the definition of NTP is in the definition of filters on $|T|_\bullet$.

As in the case of NTP, we want the map $|T|_\bullet \rightarrow M_\bullet$ in $s\mathcal{P}$ to be continuous iff there is a witness of failure of the negative requirement of item a). In case of NTP₁, it means that we define the filters on $|T|_\bullet$ by:

a subset of $|T|_\bullet(n^\leq)$ is big iff it contains a pair of incomparable vertices from each subtree of T isomorphic to ${}^{<\omega}\omega$ and all of $T^\leq(n^\leq)$, and is closed under permutations.

15. NSOP₁

Definition. A formula $\varphi(x, y)$ has SOP₁ iff there are $(a_\eta \mid \eta \in {}^{<\omega}2)$ such that

- $\forall \eta, \mu \in {}^{<\omega}2$, if $\mu \perp \eta$, then $\{\varphi(x, a_{\mu 1}), \varphi(x, a_{\eta 1})\}$ is inconsistent.
- for all branches $\sigma \in {}^{<\omega}2$, $\{\varphi(x, a_{\sigma i}) \mid i < \omega\}$ is consistent.

The reformulation is exactly the same except that we take the tree $T := {}^{<\omega}2$, and equip $|T|_\bullet(n^\leq)$ with the filter

a subset U of $|T|_\bullet(n^\leq)$ is big iff for each subtree of T isomorphic to ${}^{<\omega}2$ there is a pair $\eta, \mu \in {}^{<\omega}2$, such that $\mu \perp \eta$ and U contains all tuples containing both η and μ_1 , where μ_0 and μ_1 denote 0- and 1- descendants of μ in the subtree.

16. Rewriting No-Order-Properties

The analysis of No-Order-Properties is similar but different mathematical structures arise. Instead of an infinitely branching tree T^\leq No-Order-Properties consider the tree consisting of a single branch, i.e. an increasing sequence.

The order properties care about consistency of instances of a formula with ordered variables, hence we need to modify the filters on $|M|_\bullet$ to reflect that.

17. Rewriting NOP (no order property)

Recall a formula $\varphi(-, -)$ has NOP (no order property) iff there no sequence $(a_i)_{i \in \omega}$ such that $\varphi(a_i, a_j) \Leftrightarrow i \leq j$.

Because NOP cares about consistency of instances of a formula on tuples of ordered variables, equip $M_\bullet(n^\leq) := M^n$ with the filter of subsets containing

$$\{(a_1, \dots, a_n) \in M^n \mid i < j \ \& \ a_i \neq a_j \implies \varphi(a_i, a_j)\}$$

Then rewrite the standard definition of NOP in a form with two items representing positive and negative requirements.

Definition. A formula $\varphi(x, y)$ has NOP iff there is no sequence $(a_i)_{i \in \omega}$ such that

- for $i < j$, $\varphi(a_j, a_i)$ is false.
- for $i < j$, $\varphi(a_i, a_j)$ holds.

In analogy with NTP, think of the sequence as a tree $T := \omega^\leq$ with a single branch, and equip $\omega^\leq(n^\leq) := \text{Hom}_{\text{preorders}}(n^\leq, \omega^\leq)$, $n > 0$. with indiscrete filters. Let $|\omega|_\bullet^{\text{NOP}}(n^\leq) := \text{Hom}_{\text{Sets}}(n^\leq, |\omega|) = |\omega|^n$, $n > 0$. where a subset U of $|\omega|_\bullet^{\text{NOP}}(n^\leq) = |\omega|^n$ is large iff it is closed under permutations and

each infinite subsequence $(a_{i_j})_j$ of ω has a subsequence $(a_{i_{j_1}}, \dots, a_{i_{j_n}}) \in U$ where $j_1 > \dots > j_n$

With these definitions, $\omega^\leq \rightarrow |\omega|_\bullet^{\text{NOP}} \times M_\bullet \rightarrow \top$ defines NOP.

18. NSOP_ℓ for $\ell \geq 3$ (no strict order property)

Definition. For $\ell \geq 3$, a formula $\varphi(x, y)$ has SOP_ℓ (ℓ -strong order property) iff

- the set $\{\varphi(x_1, x_2), \dots, \varphi(x_{n-1}, x_n), \varphi(x_\ell, x_1)\}$ is inconsistent.
- there are $(a_i)_{i < \omega}$ such that $\models \varphi(a_i, a_j)$ for all $i < j$

A theory is NSOP_ℓ if no formula has SOP_ℓ.

Equivalently, NSOP_ℓ means that "item a) implies NOP".

Let $|\ell|_\bullet(n^\leq) := \text{Hom}_{\text{Sets}}(n^\leq, |\ell|) = |\ell|^n$, $n > 0$ denote the simplicial set represented by the set $\{1, 2, \dots, \ell\}$ with ℓ elements which we denote by $|\ell|$. Equip $|\ell|_\bullet(2^\leq) = |\ell|^2$ with the filter of subsets containing

$$\{(1, 2), (2, 3), \dots, (\ell - 1, \ell), (\ell, 1), (1, 1), \dots, (\ell, \ell)\},$$

and for each $n > 0$ equip $|\ell|_\bullet(n^\leq) = |\ell|^n$ with the coarsest filter such that all simplicial maps $|\ell|_\bullet(n^\leq) \rightarrow |\ell|_\bullet(2^\leq)$ are continuous, i.e. a subset is big iff it contains each element of form (i, i, \dots, i) and $(i, \dots, i, i + 1, \dots, i + 1)$, $1 \leq i \leq \ell$, and $(\ell, \dots, \ell, 1, \dots, 1)$.

Then a map $|\ell|_\bullet^{\ell\text{-cycle}} \rightarrow M_\bullet$, $i \mapsto a_i$ is continuous iff either $a_1 = \dots = a_\ell$ or $M \models \{\varphi(a_{j_1}, a_{j_2}), \dots, \varphi(a_{j_{\ell-1}}, a_{j_\ell}), \varphi(a_{j_\ell}, a_{j_1})\}$ holds for the subsequence of $(a_{j_k})_k$ of distinct elements. Hence, item a) holds iff $|\ell|_\bullet^{\ell\text{-cycle}} \rightarrow \top \times M_\bullet \rightarrow \top$. Then NSOP_{ℓ} := NSOP_{1} \vee \dots \vee NSOP_ℓ can be stated as:}}

$$|\ell|_\bullet^{\ell\text{-cycle}} \rightarrow \top \times M_\bullet \rightarrow \top \text{ implies } \omega^\leq \rightarrow |\omega|_\bullet^{\text{NOP}} \times M_\bullet \rightarrow \top$$

19. NFCP (no finite cover property)

We also note that in the category $s\mathcal{P}$ of generalised topological spaces NFCP (no finite cover property) means that M_\bullet is of infinite dimension. Recall that " M_\bullet is of finite dimension" (as a simplicial object) means that the whole of M_\bullet is fully determined by a finite piece of M_\bullet , namely there is $N > 0$ such that for each $n > N$ $M_\bullet(n^\leq)$ is the pullback of the diagram consisting of simplicial maps $M_\bullet(n^\leq) \rightarrow M_\bullet(m_1^\leq)$ and $M_\bullet(m_1^\leq) \rightarrow M_\bullet(m_2^\leq)$ where $m, m_1, m_2 < N$. In terminology of Malliaris [6, Remark 2.6] this is expressed by saying that "the characteristic sequence $\langle P_n \rangle$ of φ has finite support".

20. Stability and NIP

Both stability and NIP can be expressed in terms of indiscernible sequences. Hence, we modify the filters on $|M|_\bullet$ to reflect that: the filter on $|M|_\bullet(n^\leq) = M^n$ is generated by sets of φ -indiscernible sequences with repetitions, i.e. sequences $(a_i)_i$ such that $\varphi(a_{i_1}, \dots, a_{i_r}) \Leftrightarrow \varphi(a_{j_1}, \dots, a_{j_r})$ whenever $i_1 < \dots < i_r$, $j_1 < \dots < j_r$, all the a_{i_1}, \dots, a_{i_r} are distinct, and all the a_{j_1}, \dots, a_{j_r} are distinct. (If you feel it is more natural, you may require each subsequence with distinct elements to be part of an infinite φ -indiscernible sequence.)

This construction of M_\bullet generalises to formulas of arbitrary arity, and, moreover, for arbitrary collections of formulas.

There is a forgetful functor $s\mathcal{P} \rightarrow \text{Top}$ taking a generalised topological space into a topological space (possibly empty). For a unary formula $\varphi(-)$, it takes M_\bullet defined above into the (usual) Stone space of φ -types.

21. Each indiscernible sequence is a set.

Let $\omega_{\bullet}^{\leq, \text{cof}}(n^{\leq}) := \text{Hom}_{\text{preorders}}(n^{\leq}, \omega^{\leq}), n > 0,$
 $|\omega|_{\bullet}^{\text{cof}}(n^{\leq}) := \text{Hom}_{\text{sets}}(n^{\leq}, |\omega|) = |\omega|^n, n > 0$

equip $|\omega| = \omega_{\bullet}^{\leq, \text{cof}}(1^{\leq}) = |\omega|_{\bullet}^{\text{cof}}(1^{\leq})$ with the filter of cofinite subsets, and equip each $\omega_{\bullet}^{\leq, \text{cof}}(n^{\leq}),$ resp. $|\omega|_{\bullet}^{\text{cof}}(n^{\leq}),$ with the coarsest filter such that all simplicial maps $\omega_{\bullet}^{\leq, \text{cof}}(n^{\leq}) \rightarrow \omega_{\bullet}^{\leq, \text{cof}}(2^{\leq}),$ resp. $|\omega|_{\bullet}^{\text{cof}}(n^{\leq}) \rightarrow |\omega|_{\bullet}^{\text{cof}}(2^{\leq}),$ are continuous. Then

each (infinite) φ -indiscernible sequence is a φ -indiscernible set iff
 $\omega_{\bullet}^{\leq, \text{cof}} \rightarrow |\omega|_{\bullet}^{\text{cof}} \times M_{\bullet} \rightarrow \top$

22. Each indiscernible sequence is eventually indiscernible over any parameter

Let $M[+1]_{\bullet}(n^{\leq}) := \text{Hom}_{\text{sets}}((n+1)^{\leq}, |\omega|) = |M| \times |M|^n, n > 0,$
and equip $|M|^{n+1} = M[+1]_{\bullet}(n^{\leq})$ with the filter generated by subsets

$\{(a_0, a_1, \dots, a_n) : \text{the seq. } (a_1, \dots, a_n) \text{ is } \varphi\text{-indiscernible over } a_0\}, \varphi \text{ a formula}$

One can check that the map $M[+1]_{\bullet} \rightarrow M_{\bullet}, (a_0, a_1, \dots, a_n) \mapsto (a_1, \dots, a_n)$ forgetting the first coordinate, is continuous.

Each indiscernible sequence in M is eventually indiscernible over any parameter iff

$$\{0\}_{\bullet} \rightarrow \omega_{\bullet}^{\leq, \text{cof}} \times M[+1]_{\bullet} \rightarrow M_{\bullet}$$

where $\{0\}_{\bullet}(n^{\leq}) = \{0\}$ is equipped with the filter containing the empty set, and $\{0\}_{\bullet} \rightarrow \omega_{\bullet}^{\leq, \text{cof}}$ is the map $(0, \dots, 0) \mapsto (0, \dots, 0)$. Note that the map $\{0\}_{\bullet} \rightarrow M[+1]_{\bullet}$ picks an arbitrary element of M .

23. Speculations: limits and contractible models ?

In $s\mathcal{P}$ the notions of homotopy and limit are closely related:

the same construction in $s\mathcal{P}$ describes both

- picking a homotopy contracting a topological space (via singular complexes)
- taking a limit of a filter on a topological space,

and

- applies to an arbitrary morphism in $s\mathcal{P}$.

Can one somehow apply this construction to models ? One may perhaps hope that this construction can somehow be used to define a useful notion of a *contractible* (generalised topological space associated with a) *model*, or that the notion of a *limit of a combinatorial structure on a model*, say a tree as in NTP, is useful (i.e. taking a limit of the morphism associated with a combinatorial structure on a model, such as those arising in the lifting properties describing NTP).

Below we sketch this construction; see [9, §3] for more details. We warn the reader that our considerations here are unusually preliminary, and apologise for including them; our excuse is that we are likely unable to pursue them.

Definition ((Limit in a generalised space)). Let $\mathfrak{F}_{\bullet} : F_{\bullet} \rightarrow X_{\bullet}$ be a morphism in $s\mathcal{P}$. A morphism $x_{\bullet} : X_{\bullet} \rightarrow X_{\bullet} \circ [+1]$ is said to be a limit morphism (or simply a limit) of \mathfrak{F}_{\bullet} , or a homotopy contracting \mathfrak{F}_{\bullet} within X_{\bullet} , iff the following diagram commutes:

$$\begin{array}{ccc} & & X_{\bullet} \circ [+1] \\ & \nearrow x_{\bullet} & \downarrow \text{pr}_{2,3,\dots} \\ F_{\bullet} & \xrightarrow{\mathfrak{F}_{\bullet}} & X_{\bullet} \end{array}$$

where $[+1] : \Delta \rightarrow \Delta$ is the shift

$$n \mapsto n+1, f : n \rightarrow m \mapsto f' : n+1 \rightarrow m+1, f'(0) := 0; f'(i+1) := f(i) \text{ for } i \geq 0,$$

and $X_{\bullet} \circ [+1] \rightarrow X_{\bullet}$ is the expected map “forgetting the first coordinate”.

To recover the Bourbaki definition of a limit of a filter \mathfrak{F} on a topological space X , associate with $\mathfrak{F} \in \text{Ob } \mathcal{P}$ the simplicial set represented by X

$$F_{\bullet}(n^{\leq}) := \text{Hom}(n^{\leq}, X)$$

equip $F_{\bullet}(1^{\leq}) = X$ with \mathfrak{F} , and equip each $F_{\bullet}(n^{\leq}) := \text{Hom}(n^{\leq}, X)$ with the finest filter such that the diagonal map $F_{\bullet}(1^{\leq}) \rightarrow F_{\bullet}(n^{\leq})$ is continuous. A verification shows that (possibly discontinuous) liftings correspond to points of X (indeed, as simplicial sets, F_{\bullet} is connected, and $X_{\bullet} \circ [+1]$ is the disjoint union of connected component which are copies of X_{\bullet} parametrised by X), and the continuity requirement means precisely that they are limit points.

Example (Limit of the tree in NTP). Let $(a_s \mid \emptyset \neq s \in {}^{<\omega}\omega)$ be a tree of parameters such that no subtree (of the same shape) is a counterexample to NTP of a formula $\varphi(x, y)$, and all branches are φ -consistent; recall this is the same as a map $|T|_{\bullet}^{\text{NTP}} \rightarrow M_{\bullet}^{\varphi\text{-NTP}}$ in $s\mathcal{P}$. A limit or contracting homotopy of the tree $(a_s \mid \emptyset \neq s \in {}^{<\omega}\omega)$ is an element $a \in M$ such that (a) no subtree (of the same shape) of $(a_s \mid \emptyset \neq s \in {}^{<\omega}\omega)$ is a counterexample to NTP of formula $\varphi(x, a) \& \varphi(x, y)$, and (b) $\varphi(x, a), \varphi(x, a_{s_1}), \dots, \varphi(x, a_{s_n})$ is consistent whenever $n > 0$ and $s_1 < \dots < s_n$ lie on the same branch of ${}^{<\omega}\omega$.

Thus, “each NTP tree of parameters is contractible” would mean that for each tree of parameters on ${}^{<\omega}\omega$ (i) with $\varphi(x, y)$ -consistent branches and (ii) having no subtree of the same shape ${}^{<\omega}\omega$ with TP for $\varphi(x, y)$, there is some $a \in M$ such that it has the same properties with respect to $\varphi(x, a) \& \varphi(x, y)$.

To explain this example, first note that, as simplicial sets (i.e. if we ignore the filters), $M_{\bullet}^{\varphi\text{-NTP}} \circ [+1]$ is the disjoint union of connected component which are copies of $M_{\bullet}^{\varphi\text{-NTP}}$ parametrised by elements of M , and that $|T|_{\bullet}$ is connected. Thus to give a map of simplicial sets $|T|_{\bullet} \rightarrow M_{\bullet}^{\varphi\text{-NTP}}$ is the same as to pick a point $a_0 \in M$. The continuity requirement on $|T|_{\bullet}^{\text{NTP}} \rightarrow M_{\bullet}^{\varphi\text{-NTP}}$ means precisely items (a) and (b).

Can localisation and persistence of configurations in $M_{\bullet}^{\varphi\text{-NTP}}$ of Malliaris [7,8] be rephrased in terms of $s\mathcal{P}$ limits and homotopies of this kind ? Can NIP be?

24. Contracting homotopy as a limit

Now let F and X denote “nice” topological spaces.

Remark (Homotopy as limit of singular complexes). A map $h_0 : F \rightarrow X$ is *contractible*, i.e. it factors through the cone of F as $F \xrightarrow{x \mapsto (x,0)} F \times [0,1] / F \times \{1\} \xrightarrow{h} X$, iff in $s\text{Sets}$ or $s\mathcal{P}$ the map $\text{sing } F_{\bullet} \rightarrow \text{sing } X_{\bullet}$ of singular complexes has a limit i.e. there is a commutative diagram

$$\begin{array}{ccc} & & \text{sing } X_{\bullet} \circ [+1] \\ & \nearrow h_{\bullet} & \downarrow \text{pr}_{2,3,\dots} \\ \text{sing } F_{\bullet} & \longrightarrow & \text{sing } X_{\bullet} \end{array}$$

Recall that the singular complex is defined using simplices $\Delta^n = \text{Hom}_{\text{preorders}}([0,1]^{\leq}, (n+1)^{\leq})$ as “test spaces”:

$$\text{sing } F_{\bullet}((n+1)^{\leq}) := \text{Hom}_{\text{Top}}(\Delta^n, F),$$

$$\text{sing } X_{\bullet}((n+1)^{\leq}) := \text{Hom}_{\text{Top}}(\Delta^n, X),$$

$$\text{sing } X_{\bullet} \circ [+1]((n+1)^{\leq}) = \text{Hom}_{\text{Top}}(\Delta^n \times [0,1] / \Delta^n \times \{1\}, X)$$

where $n \geq 0$ and $\Delta^n \times [0,1] / \Delta^n \times \{1\}$ is the cone of n -simplex Δ^n .

To define a limit(=lifting) h_{\bullet} , take each $\delta : \Delta^n \rightarrow F$ in $F_{\bullet}((n+1)^{\leq})$ to $h_{\bullet}(\delta) : \Delta^n \times [0,1] / \Delta^n \times \{1\} \rightarrow X$ in $X_{\bullet}((n+2)^{\leq})$ defined by

$$h_{\bullet}(\delta)(x, t) := h(\delta(x), t).$$

To see the other direction, note that $h_{\bullet} : F_{\bullet} \rightarrow X_{\bullet} \circ [+1]$ takes a singular simplex $\delta : \Delta^n \rightarrow F$ into $h_{\bullet}(\delta) : \Delta^{n+1} = \Delta^n \times [0,1] / \Delta^n \times \{1\} \rightarrow X$ such that $\delta \circ h_0 = h_{\bullet}(\delta)|_{\Delta^n \times \{0\}}$, i.e. each $\delta : \Delta^n \rightarrow F \rightarrow X$ factors through the cone of Δ^n . A verification using functoriality shows that the same factorisation holds for $\mathbb{S}^n = \partial \Delta^{n+1}$, which means exactly that h_0 is weakly contractible, and for “nice” topological spaces contractible and weakly contractible are equivalent.

Remark (Homotopy as limit of the fibre). A map $h : F \times [0,1] / F \times \{1\} \rightarrow X$ continuous in a neighbourhood of “the top of the cone” point $F \times \{1\}$ is the same as a map in $s\mathcal{P}$

$$F_{\bullet} \times ([0,1]_1)_{\bullet} \rightarrow X_{\bullet} \circ [+1]$$

$$(x_1, \dots, x_n, t_1, \dots, t_n) \mapsto (h(x_1, 1), h(x_1, t_1), h(x_2, t_2), \dots, h(x_n, t_n))$$

where $[0,1]_1$ denotes the interval $[0,1]$ equipped with the filter of neighbourhoods of point 1, and F_{\bullet}, X_{\bullet} denote the generalised topological spaces corresponding to F and X .

In other words, $s\mathcal{P}$ can express infinitesimally/“sufficiently” short homotopies: in an expressive language, we may say that $h_t : F \rightarrow X, t \in [0,1]$, converges at 0 iff there is $\varepsilon > 0$ such that $h|_{[0,\varepsilon]}$ is a homotopy contracting F in X .

25. Homotopy theory for a model ?

A standard way to modify the definition of $\text{sing} : \text{Top} \rightarrow s\text{Sets}$ (or $\text{nerve} : \text{Cat} \rightarrow s\text{Sets}$) is to use instead of simplices another cosimplicial “test” space, i.e. take a functor $\mathfrak{a}^* : \text{Non-emptyFiniteLinearOrders} \rightarrow s\mathcal{P}$ and set

$$\text{sing } X_{\bullet}(n^{\leq}) := \text{Hom}(\mathfrak{a}^*(n^{\leq}), X)$$

where Hom may mean a space (rather than merely a set) of maps, e.g. internal hom or as defined in [8]. This suggest that we find a cosimplicial “test” object in $s\mathcal{P}$ such that $\text{Hom}(\mathfrak{a}^*(n^{\leq}), M_{\bullet})$ is meaningful in model theory, perhaps by finding first a sequence of interesting combinatorial structures.

Alternatively, do something like what we did with NTP: find a definition in model theory that has the pattern of the diagram defining limit in $s\mathcal{P}$. Say, you'd like to see (force?) that NTP_k has pattern of $\pi_n(X) = 0$ e.g. that $T_{\bullet}^{\leq} \rightarrow |T|_{\bullet}^{\text{NTP}_k} \times M_{\bullet}^{\varphi\text{-NTP}} \rightarrow \top$ has the pattern of $\partial \Delta^{n+1} \rightarrow \Delta^{n+1} \times X \rightarrow \top$, defining $\pi_n(X) = 0$ in $s\text{Sets}$. Hence, you'd want to see how monotone maps $(n+2)^{\leq} \rightarrow (n+2)^{\leq}$ would act on $|T|_{\bullet}^{\text{NTP}_k}$ (but not T_{\bullet}^{\leq}). You'd also want this action if you want $|T|_{\bullet}^{\text{NTP}_k}$ to be part of a cosimplicial object used to define sing_{\bullet} . (Note that in topology there is no *natural* action of these maps on \mathbb{D}^{n+1} occurring in $\mathbb{S}^n \rightarrow \mathbb{D}^{n+1} \times X \rightarrow \top$ defining $\pi_n(X) = 0$: you need to pick an arbitrary homeomorphism of ball \mathbb{D}^{n+1} with the standard $(n+1)$ -simplex.)

Acknowledgements.

Without help, insight, and efforts of M.Bays and A.Hasson our reformulations would never have appeared. V.Sonsilo pointed out that the $s\mathcal{P}$ reformulation of limit reminds of homotopy, and was willing to bear with our category-theoretic questions. Will Johnson pointed out many inaccuracies in our earlier treatment of no-order-properties and stability. See [9] for history and more acknowledgements. The first page is a (corrected) poster prepared for SCS'2022. The authors are likely in no position to develop ideas presented.

References.

1. A.Pillay, S.Starchenko. Remarks on Tao's algebraic regularity lemma.. 2013.
2. P.Simon. A note on “regularity lemma for distal structures”. 2015.
3. An overview of generalised topological spaces. ncatlab.com/show/situs
4. Lifting property. en.wikipedia.org/wiki/Lifting_property ncatlab.com/show/lift
5. K.Tent, M.Ziegler. A course in model theory.
6. M.Malliaris. Edge distribution and density in the characteristic sequence, 2010.
7. M.Malliaris. The characteristic sequence of a first-order formula, 2010,
8. M.Gavrilovich, K.Pimenov. Geometric realisation as the Skorokhod semi-continuous path space endofunctor.
9. An informal exposition “reading off” the definition of the lifting property and that of $s\mathcal{P}$ from (Bourbaki, General Topology): mishap.sdf.org/ yet_another_not_an_obfuscated_story.pdf

Contact: gavrilovich@ihes.fr text chat: t.me/McVlr

During poster room hours at P&S'60: voice chat mishap.sdf.org/gathertown